THE SINGULARITIES OF A_g^P

J. ARMSTRONG AND N. I. SHEPHERD-BARRON

arXiv:1604.05943v1 [math.AG] 20 Apr 2016

Let $\mathcal{A}_g$ denote the stack, and $A_g = [\mathcal{A}_g]$ the coarse moduli space, of principally polarized abelian g-folds and $\mathcal{A}_g^P$, $A_g^P = [\mathcal{A}_g^P]$ their perfect cone compactifications; these are particular toroidal compactifications.

The point of this paper is to complete the proof in [SB] of a result that is suggested by Tai's paper [T], that in characteristic zero the singularities of A_g^P are canonical if $g \geq 5$ and terminal if $g \geq 6$. However, Hulek and Sankaran have pointed out that the argument given in [SB] is incomplete, because Proposition 3.2 of [SB] is not strong enough. We complete it by replacing Proposition 3.2 of *loc. c it.* with Theorem 1.1 below.

1 The proof

We recall the RST criterion for the geometric quotient $[\mathcal{X}]$ of a smooth Deligne–Mumford stack $\mathcal{X}$ over a field of characteristic zero to have canonical or terminal singularities. We phrase it in terms of the action of a finite group G acting on a smooth variety Y. First, let $\{x\}$ denote the fractional part of a real number x. Then, for any $s \in G - \{1\}$, define $\lambda_{Y,P}(s) = \sum \arg(\zeta)/2\pi$, where the sum is over the eigenvalues ζ of s on the tangent space $T_{Y,P}$.

(RST) G acts freely in codimension one in a neighbourhood of P and $[Y/G]$ has canonical (resp., terminal) singularities at the image of P if and only if $\lambda_{Y,P}(s) \geq 1$ (resp., $\lambda_{Y,P}(s) > 1$) for every $s \in G - \{1\}$ and every choice of ζ.

Theorem 1.1 *The singularities of A_g^P are canonical if $g = 5$, and terminal if $g \geq 6$.*

PROOF: The problem reduces to this. Suppose that $h, r \geq 0$, $g = h + r$, C is a principally polarized abelian h-fold, $W = H^0(C, \Omega_C^1)^\vee$, $\Lambda = \mathbb{Z}^r$, $V = \operatorname{Sym}^2 W \oplus (W \otimes \Lambda)$ and $\mathbb{T}$ is the torus with $\mathbb{X}_*(\mathbb{T}) = B(\Lambda)$, the group of symmetric $\mathbb{Z}$-valued bilinear forms on Λ. There is a torus embedding $\mathbb{T} \hookrightarrow X$ with terminal singularities and a $\mathbb{T}$-bundle $\mathcal{T} \to V$ such that locally $\mathcal{A}_g^P = \mathcal{X}/G$ for a finite group $G = \operatorname{Stab}_{\mathcal{X}}(P)$, where $\mathcal{X} = \mathcal{T} \times^{\mathbb{T}} X \to V$ is a G-equivariant X-bundle and $P \in \mathcal{X}$. This is compatible with a $\mathbb{Z}$-linear action of G on Λ and an action of G on C, so that $\mathbb{T} \hookrightarrow X$ is G-equivariant and G acts linearly on V. In terms of co-ordinates, choose a $\mathbb{Z}$-basis $(x_1, \dots, x_r)$ of Λ and write $Z_{pq} = \exp 2\pi i(x_p \odot x_q)$. Then locally $\mathcal{O}_{\widetilde{\mathcal{X}}} = \mathcal{O}_V[Y_m]$ for some monomials Y_m in the Z_{pq} and the action of an element $s \in G$ is given by $s^*Y_m = \Phi_{s,m} W_{s(m)}$, where $\Phi_m \in \mathcal{O}_V^{hol}$, Φ_m does not vanish at 0 and $W_{s(m)}$ is a monomial in the Z_{pq}.

Lemma 1.2 *If $h \geq 1$, then G acts effectively on V.*

PROOF: The only elements of $GL(W) \times GL(\Lambda)$ that act trivially on V are ± 1. □

The next proposition slightly extends one of Tai's. We omit the proof, but it follows his almost exactly: reduce to the case where the order of s divides 12 and then check a finite list. We executed this check by writing a routine in Mathematica.

Proposition 1.3 *Assume that $g \geq 5$, that $h \geq 1$ and that $s \in G - \{1\}$.*

(1) If $s|_W \neq \pm 1$, then $\lambda_{\mathrm{Sym}^2 W}(s) \geq 1$ if $h = 5$ and $\lambda_{\mathrm{Sym}^2 W}(s) > 1$ if $h \geq 6$.

(2) If $s|_V$ is not of order 2, then $\lambda_V(s) \geq 1$.

(3) If $\lambda_V(s) < 1$ then $h = 1$, $s|_W = -1$, $s|_{\Lambda_{\mathbb{Q}}} = (1, (-1)^{r-1})$ and $\lambda_V(s) = 1/2$.

In particular, taking $g = h$ shows that the singularities of the interior A_g are terminal if $g \geq 6$ and canonical if $g = 5$.

For the proof of the theorem, choose a $\mathbb{T} \rtimes G$-equivariant resolution $\widetilde{X} \to X$, and put $\widetilde{\mathcal{X}} = \mathcal{T} \times^{\mathbb{T}} \widetilde{X}$. If s acts trivially on some divisor E in $\widetilde{\mathcal{X}}$, then E is in the boundary $\partial\widetilde{\mathcal{X}} = \widetilde{\mathcal{X}} - \mathcal{T}$. Each boundary divisor dominates V, since $\widetilde{\mathcal{X}} \to V$ is a fibre bundle, and then s acts trivially on V. This is false, so that G acts freely in codimension one on $\widetilde{\mathcal{X}}$.

For some $q \geq 1$ there is a G-invariant generator σ of the sheaf $\mathcal{O}(qK_{\mathcal{X}})$. Then σ is also a generator of $\mathcal{O}(qK_{[\mathcal{X}/G]})$. Since $\mathcal{X}$ is smooth, σ vanishes along all exceptional divisors in $\widetilde{\mathcal{X}}$, so that σ also vanishes along all exceptional divisors in $[\widetilde{\mathcal{X}}/G]$. So it is enough to deal with the singularities of $[\widetilde{\mathcal{X}}/G]$.

Suppose that $h \geq 1$ and $g \geq 6$. Assume that $\lambda_{\widetilde{\mathcal{X}},P}(s) \leq 1$. Note that $\lambda_{\widetilde{\mathcal{X}},P}(s) \geq \lambda_V(s) + \lambda_{\widetilde{\mathcal{X}}/V,P}(s)$, from the definition of λ, and that P lies in $\partial\widetilde{\mathcal{X}}$, since $\lambda_{\mathcal{A}_g,Q} > 1$ for all points Q of $\mathcal{A}_g$. If $\lambda_{\widetilde{\mathcal{X}}/V,P}(s) = 0$ then s acts trivially on the relative tangent space $T_{\widetilde{\mathcal{X}}/V,P}$, so that a suitable point $Q \in \mathcal{T}$ that is near P would give $Q \in \mathcal{A}_g$ with $\lambda_{\mathcal{A}_g,Q}(s) = \lambda_{V,P}(s) \leq 1$. So $\lambda_{\widetilde{\mathcal{X}}/V,P}(s) > 0$, $\lambda_V(s) < 1$, and then s is of order 2, $\lambda_V(s) = 1/2$, $h = 1$ and $r \geq 5$. Moreover, $\lambda_{\widetilde{\mathcal{X}}/V,P}(s) = 1/2$.

There is an s-equivariant factorization $\widetilde{\mathcal{X}} \xrightarrow{\alpha} \mathcal{Y} \xrightarrow{\beta} V$ of $\widetilde{\mathcal{X}} \to V$, where α is a vector bundle and β is a torus bundle. The algebra $\mathcal{O}_{\mathcal{Y}}$ is generated as an $\mathcal{O}_V$-algebra by those monomials Y_m that are invertible at P.

If $\lambda_{\widetilde{\mathcal{X}}/\mathcal{Y},P}(s) = 0$, then we could, as before, find $Q \in \mathcal{A}_g$ with $\lambda_{\mathcal{A}_g,Q}(s) = 1$, contradiction. So $\lambda_{\widetilde{\mathcal{X}}/\mathcal{Y},P}(s) = 1/2$ and $\lambda_{\mathcal{Y}/V,P}(s) = 0$. We have $\mathcal{O}_{\widetilde{\mathcal{X}}} = \mathcal{O}_{\mathcal{Y}}[Y_1, \ldots, Y_n]$ and $s^*Y_i = \Psi_i Y_{s(i)}$, with $\Psi_i \in \mathcal{O}_{\mathcal{Y}}^*$. Since $\lambda_{\widetilde{\mathcal{X}}/\mathcal{Y},P}(s) = 1/2$, the trace of s acting on $T_{\widetilde{\mathcal{X}}/\mathcal{Y},P}$ equals $\dim T_{\widetilde{\mathcal{X}}/\mathcal{Y},P} - 2$, so that there at most two indices i for which $s(i) \neq i$. Since also $\lambda_{\mathcal{Y}/V,P}(s) = 0$, there is a $\mathbb{Q}$-basis of $B(\Lambda_{\mathbb{Q}})$ of which s fixes at least $\binom{r+1}{2} - 2$ elements. But $s|_{\Lambda\otimes\mathbb{Q}} = (1, (-1)^{r-1})$ and $r \geq 5$, so this is impossible and we have contradicted the assumption that $\lambda_{\widetilde{\mathcal{X}},P} \leq 1$.

The proof that A_5^P has canonical singularities when $h \geq 1$ is the same.

Finally, if $h = 0$, then we use a theorem of Snurnikov [Sn]. Since it remains unpublished, we include a proof.

Proposition 1.4 *Suppose $\mathbb{T} \hookrightarrow X$ is a G-equivariant torus embedding, $U \subset X$ a G-invariant neighbourhood of $X - \mathbb{T}$, G acts freely in codimension one on $U \cap \mathbb{T}$ and that U and $[(U \cap \mathbb{T})/G]$ have terminal (canonical) singularities. Then G acts freely in codimension one on U and $[U/G]$ has terminal (canonical) singularities.*

PROOF: Let $\pi : \widetilde{X} \to X$ be a $\mathbb{T} \rtimes G$-equivariant resolution. As before, it is enough to show that the G-action on $\widetilde{U} = \pi^{-1}(U)$ satisfies (RST).

If $s \in G$ and $s(P) = P$, then, locally near P, $\widetilde{X}$ is an s-equivariant vector bundle $q : \widetilde{X} = \mathbf{Spec}\,\mathbb{C}[x, y_j^{\pm}] \to \mathbb{T}_1 = \mathbf{Spec}\,\mathbb{C}[y_j^{\pm}]$; the y_j are those characters of $\mathbb{T}$ that are invertible at P and $s^* x_i = \phi_i x_{s(i)}$, where ϕ_i is a monomial in the y_j. Let R denote the origin in the fibre $F = q^{-1}(q(P))$; then $\lambda_{\widetilde{U},P}(s) = \lambda_{\widetilde{U},R}(s)$, since $s \mapsto \lambda(s)$ is constant on the line in F through R and P.

So we can assume that $P = R$ and that $\widetilde{X} \neq \mathbb{T}_1$. Then the locus $x_i = t$, $y_j^{\pm} = 1$ for all i, j is a line L of s-fixed points in $\widetilde{X}$ that passes through P and meets $\widetilde{U} \cap \mathbb{T} = U \cap \mathbb{T}$. Suppose $Q \in U \cap L$ is general; then $\lambda_{\widetilde{X},P}(s) = \lambda_{U \cap \mathbb{T},Q}(s)$. □

This completes the proof of the theorem. □

References

[SB] N.I. Shepherd-Barron, *Perfect forms and the moduli space of abelian varieties*, Invent. Math. (2006).

[S] V.S. Snurnikov, Ph.D. thesis, Cambridge, 2002.

[T] Y.-S. Tai, *On the Kodaira dimension of the moduli space of abelian varieties*, Invent. Math. **68** (1982), 425-439.

MATH. DEPT., KING'S COLLEGE LONDON, STRAND, WC2R 2LS, UK
E-mail address: john.armstrong@kcl.ac.uk, nicholas.shepherd-barron@kcl.ac.uk